Government of Saskatchewan

# Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 06, June 2, 2016

**For the Period May 24 to 30, 2016**

Seeding of the 2016 crop is almost complete. Ninety four per cent of the crop is now seeded, according to Saskatchewan Agriculture's weekly Crop Report. Cooler temperatures and rain during the past week have slowed field operations in some areas of the province; however, seeding progress is still ahead of the five-year (2011-2015) average of 77 per cent for this time of year. Many producers have completed seeding operations and are now starting in-crop herbicide applications.

Ninety-six per cent of the crop is seeded in the southeast, 95 per cent in the west central and northwest regions, 94 per cent in the northeast, 93 per cent in the southwest and 91 per cent in the east central region.

Rain was fairly general across the province and heavy downpours have been reported in some areas, particularly in southern regions. Precipitation ranged from zero to 95 mm. Some areas will still need rain in the coming weeks to help crops emerge and establish. Topsoil moisture conditions on cropland are rated as eight per cent surplus, 86 per cent adequate and six per cent short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 84 per cent adequate, eight per cent short and two per cent very short.

Overall, crops that have emerged are in good to excellent condition, although there has been some damage caused by flea beetles, cutworms, localized flooding and lack of moisture.

Producers are busy completing seeding operations and controlling weeds and insects.

**One year ago**
Ninety-seven per cent of the crop had been seeded. Heavy frost in some areas caused crop damage. Due to dry conditions emergence and germination was of concern in some areas of the province.

Follow the 2016 Crop Report on Twitter @SKAgriculture

| Seeding Progress in SK Per cent seeded All Crops | |
|---|---|
| May 30, 2016 | **94** |
| June 1, 2015 | 97 |
| June 2, 2014 | 78 |
| May 27, 2013 | 67 |
| May 28, 2012 | 74 |
| May 30, 2011 | 70 |
| **5 year avg. (2011-2015)** | **77** |
| **10 year avg. (2006-2015)** | **80** |

| Saskatchewan Seeding Progress May 30, 2016 % Seeded | |
|---|---|
| Spring wheat | 95 |
| Durum | 94 |
| Barley | 90 |
| Oats | 87 |
| Canary seed | 86 |
| Flax | 86 |
| Canola | 94 |
| Mustard | 93 |
| Soybeans | 92 |
| Lentils | 98 |
| Field peas | 98 |
| Chickpeas | 97 |

For further information, contact Daphne Cruise, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3587, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

**Southeastern Saskatchewan (Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas; Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas; Crop District 3ASE – Radville and Lake Alma areas)**

Seeding is 96 per cent complete in the region. The five-year (2011-2015) seeding average for this time of year is 68 per cent. Significant amounts of rain throughout the week slowed field operations, with most areas reporting over 35 mm of rain. The rain was welcome, as conditions were dry for emerging crops and hay and pasture land. Although, excess rain over a short amount of time in some areas has caused issues such as flooding and has affected field access.

Rainfall recorded ranged from 20 mm (Vibank area) to 95 mm (Moosomin area). Other areas receiving larger amounts of rain included Lampman (61 mm), Weyburn (47 mm) and Radville (56 mm). The Moose Jaw area holds the regional record for greatest amount of rainfall received since April 1 (155 mm). The Vibank area has the least accumulated rain in the region (40 mm) since April 1.

Topsoil moisture conditions have improved over last week, with 16 per cent rated as surplus, 82 per cent adequate and two per cent short on cropland. Hay land and pasture topsoil moisture is rated as 13 per cent surplus, 84 per cent adequate and three per cent short. Crop District 1A is reporting 30 per cent surplus topsoil moisture on crop land.

Overall, emerging crops are in good condition. The majority of crop damage this week was due to flooding, cutworm and fleabeetle damage. There are also many reports of tent caterpillars damaging trees in the region. In-crop herbicide applications are just beginning.

Farmers are busy finishing up seeding, rolling pulse crops, controlling weeds and moving cattle.

**Southwestern Saskatchewan (Crop District 3ASW – Coronach, Assiniboia and Ogema areas; Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas; Crop District 3B – Kyle, Swift Current , Shaunavon and Ponteix areas; Crop District 4 – Consul, Maple Creek and Leader areas)**

Farmers in the region have almost wrapped up seeding, with 93 per cent complete. The five-year (2011-2015) seeding average for this time of year is 83 per cent.

Rainfall in the region was general and ranged from 10 mm (Climax area) to 68 mm (Tyner area). The Limerick area received 51 mm, the Eyebrow and Gull Lake areas 45 mm, the Hazenmore area 54 mm and the Leader area 53 mm. The Shaunavon area holds the regional, and provincial, record for greatest amount of rainfall received since April 1 (216 mm). Cropland topsoil moisture is rated as eight per cent surplus, 91 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as five per cent surplus, 92 per cent adequate and three per cent short. Crop district 3ASW is reporting 20 per cent surplus of topsoil moisture on crop and hay land.

Emerged crops are in good to excellent condition, although many fields remain wet from recent rain. The rain is welcomed after a year of very dry conditions in 2015. However, due to significant rainfall, there have been reports of localized flooding causing crop damage in some areas.

Farmers are busy completing seeding, rolling pulses and controlling weeds.

**East-Central Saskatchewan (Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas; Crop District 6A – Lumsden, Craik, Watrous and Clavet areas)**

Seeding operations are continuing in the region. Ninety-one per cent of the crop is now seeded. Some areas were able to continue with field work between rain showers. The five-year (2011-2015) seeding average for this time of year is 69 per cent.

Rainfall recorded ranged from 1 mm (Raymore area) to 46 mm (Holdfast area). Crop region 5B reported the least amount of rainfall, with most areas receiving less than 10 mm. The Bethune area holds the regional record for greatest amount of rainfall received since April 1 (124 mm). Cropland topsoil moisture conditions are rated as four per cent surplus, 88 per cent adequate, seven per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as three per cent surplus, 88 per cent adequate, seven per cent short and two per cent very short. Crop district 5B is reporting 13 per cent short on crop land topsoil moisture.

Emerged crops are in relatively good to excellent condition, although damage this week was caused by flea beetle, cutworm and flooding.

Farmers are busy seeding, controlling weeds and scouting fields for insects.

**West-Central Saskatchewan (Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas; Crop District 7A – Rosetown, Kindersley, Eston, Major; Crop District 7B - Kerrobert, Macklin, Wilkie and Biggar areas)**

Farmers in the region have 95 per cent of the 2016 crop seeded. The five-year (2011-2015) seeding average for this time of year is 83 per cent.

Some areas of the region received much-needed moisture this past week. Rainfall in the region ranged from 5 mm (Outlook and Cando areas) to 51 mm (Tramping Lake area). The Marengo area also holds the regional record for greatest amount of rainfall received since April 1 (122 mm). The Rosthern area has totalled 27 mm since April 1.

Topsoil moisture conditions have greatly improved, thanks to the recent rain. Topsoil moisture conditions on cropland are rated as eight per cent surplus, 85 per cent adequate, six per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as six per cent surplus, 78 per cent adequate, 12 per cent short and four per cent very short. At this time, Crop District 6B

is reporting that 12 per cent of cropland and hay land and pasture acres remain short topsoil moisture.

Overall, emerged crops are in good condition. The recent moisture will help to germinate crops evenly and get them off to a good start. There a few reports of cutworm and flea beetle causing damage in canola and lentil crops.

Farmers are busy completing seeding, controlling weeds, rolling pulses and scouting for insects.

**Northeastern Saskatchewan (Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas; Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas)**

The northeastern region has 94 per cent of the crop seeded. Although the region reported rain, most were able to keep to the fields between small shower events. The five-year (2011-2015) seeding average for this time of year is 80 per cent.

Rainfall ranged from two mm (Kinistino area) to 31 mm (Alvena area). The Nipawin area also holds the regional record for greatest amount of rainfall received since April 1 (106 mm). The Humboldt area has recorded 28 mm since April 1.

Cropland topsoil moisture conditions are rated as four per cent surplus, 88 per cent adequate, eight per cent short and one per cent very short. Hay land and pasture topsoil moisture is rated as two per cent surplus, 86 per cent adequate and 12 per cent short. Crop District 8A is reporting that 12 per cent of cropland and 14 per cent of hay land and pasture are short on topsoil moisture at this time.

Crops are emerging and are in relatively good condition. There were a few reports of flea beetle damage in canola in the Lake Lenore area.

Farmers are busy seeding, controlling weeds and scouting for insects.

**Northwestern Saskatchewan (Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas; Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas)**

Seeding continued to rapidly progress during the week, with 95 per cent now seeded. The five-year (2011-2015) seeding average for this time of year is 84 per cent. Although rain was welcomed in the region last week, some areas remain dry and will need additional moisture in the coming weeks.

Rainfall ranged from 2 mm (Debden area) to 47 mm (Lloydminster area). The Frenchman Butte area holds the regional record for greatest amount of rainfall received since April 1 (76 mm). The Duck Lake area has recorded 18 mm since April 1.

Topsoil moisture conditions have improved in the region, although some areas will need more rain in the coming weeks. Cropland topsoil moisture is rated as six per cent surplus, 75 per cent adequate and 19 per cent short. Hay land and pasture topsoil moisture is rated as three per cent surplus, 67 per cent adequate, 22 per cent short and eight per cent very short.

Crop emergence is generally good with the recent moisture, although germination is spotty due to the dry field conditions in some areas. Hail was reported in the Barthel and Turtleford areas, though damage is expected to be minimal, as most crops had not emerged yet.

Farmers are busy completing seeding, moving cattle and controlling weeds.

# Crop Conditions - June 2, 2016

| Provincial | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 23 | 49 | 22 | 30 | 15 | 19 | 27 |
| % good | 67 | 47 | 72 | 65 | 77 | 73 | 64 |
| % fair | 9 | 3 | 6 | 5 | 8 | 8 | 9 |
| % poor | 1 | 1 | 0 | 0 | 0 | 0 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 20 | 20 | 22 | 5 | 34 | 37 | 16 |
| % good | 73 | 70 | 76 | 85 | 61 | 58 | 78 |
| % fair | 7 | 9 | 2 | 9 | 5 | 5 | 6 |
| % poor | 0 | 1 | 0 | 1 | 0 | 0 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| Southeast | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 10 | 19 | 22 | 33 | 11 | 19 | 44 |
| % good | 70 | 70 | 65 | 58 | 79 | 67 | 47 |
| % fair | 20 | 11 | 12 | 9 | 10 | 13 | 9 |
| % poor | 0 | 0 | 1 | 0 | 0 | 1 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 21 | 24 | 9 | 3 | 22 | 35 | N/A |
| % good | 68 | 60 | 86 | 86 | 70 | 57 | N/A |
| % fair | 11 | 15 | 5 | 10 | 7 | 8 | N/A |
| % poor | 0 | 1 | 0 | 1 | 1 | 0 | N/A |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | N/A |

| Southwest | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 27 | 63 | 29 | 31 | 36 | 31 | 9 |
| % good | 68 | 35 | 68 | 65 | 60 | 64 | 75 |
| % fair | 3 | 2 | 3 | 4 | 3 | 4 | 11 |
| % poor | 2 | 0 | 0 | 0 | 1 | 1 | 5 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 19 | 19 | 26 | 3 | 43 | 36 | 15 |
| % good | 78 | 76 | 74 | 87 | 55 | 61 | 78 |
| % fair | 3 | 4 | 0 | 10 | 2 | 3 | 7 |
| % poor | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| East-central | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 38 | 15 | 16 | 29 | 12 | 12 | 0 |
| % good | 59 | 85 | 77 | 71 | 74 | 77 | 78 |
| % fair | 2 | 0 | 7 | 0 | 14 | 11 | 22 |
| % poor | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

| East-central (continued) | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 18 | 13 | N/A | 8 | 24 | 29 | N/A |
| % good | 78 | 73 | N/A | 85 | 74 | 69 | N/A |
| % fair | 4 | 14 | N/A | 7 | 2 | 2 | N/A |
| % poor | 0 | 0 | N/A | 0 | 0 | 0 | N/A |
| % very poor | 0 | 0 | N/A | 0 | 0 | 0 | N/A |

| West-central | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | 4 | 33 | 24 | 25 | 20 | 27 | 25 |
| % good | 96 | 67 | 69 | 72 | 66 | 67 | 73 |
| % fair | 0 | 0 | 7 | 3 | 14 | 6 | 2 |
| % poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 30 | 24 | 24 | 35 | 36 | 41 | N/A |
| % good | 67 | 67 | 67 | 58 | 52 | 53 | N/A |
| % fair | 3 | 8 | 8 | 7 | 12 | 6 | N/A |
| % poor | 0 | 1 | 1 | 0 | 0 | 0 | N/A |
| % very poor | 0 | 0 | 0 | 0 | 0 | 0 | N/A |

| Northeast | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | N/A | N/A | 20 | 88 | 19 | 22 | 12 |
| % good | N/A | N/A | 78 | 12 | 76 | 69 | 63 |
| % fair | N/A | N/A | 2 | 0 | 5 | 9 | 25 |
| % poor | N/A | N/A | 0 | 0 | 0 | 0 | 0 |
| % very poor | N/A | N/A | 0 | 0 | 0 | 0 | 0 |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 15 | 27 | N/A | N/A | 39 | N/A | N/A |
| % good | 70 | 71 | N/A | N/A | 59 | N/A | N/A |
| % fair | 15 | 2 | N/A | N/A | 2 | N/A | N/A |
| % poor | 0 | 0 | N/A | N/A | 0 | N/A | N/A |
| % very poor | 0 | 0 | N/A | N/A | 0 | N/A | N/A |

| Northwest | | | | | | | |
|---|---|---|---|---|---|---|---|
| | Winter wheat | Fall rye | Spring wheat | Durum | Oat | Barley | Canaryseed |
| % excellent | N/A | N/A | 23 | N/A | 10 | 9 | N/A |
| % good | N/A | N/A | 76 | N/A | 89 | 91 | N/A |
| % fair | N/A | N/A | 1 | N/A | 1 | 0 | N/A |
| % poor | N/A | N/A | 0 | N/A | 0 | 0 | N/A |
| % very poor | N/A | N/A | 0 | N/A | 0 | 0 | N/A |
| | Flax | Canola | Mustard | Soybean | Pea | Lentil | Chickpea |
| % excellent | 40 | 16 | N/A | N/A | 18 | N/A | N/A |
| % good | 60 | 78 | N/A | N/A | 80 | N/A | N/A |
| % fair | 0 | 6 | N/A | N/A | 2 | N/A | N/A |
| % poor | 0 | 0 | N/A | N/A | 0 | N/A | N/A |
| % very poor | 0 | 0 | N/A | N/A | 0 | N/A | N/A |

# Crop Districts and Rural Municipalities in Saskatchewan




Government of Saskatchewan

0 25 50 100 150 200
Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Crop Districts - Saskatchewan Ministry of Agriculture

Geomatic Services, Ministry of Agriculture June 10, 2014

© 2014 Government of Saskatchewan

# Weekly Rainfall

## from May 24 to May 30, 2016




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture June 1, 2016

© 2016 Government of Saskatchewan

# Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period May 24 to 30, 2016

| Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr |
|---|---|---|---|---|
| 1A | 2 | Mount Pleasant | 41 | 95 |
| | 3 | Enniskillen | 37 | 85 |
| | 33 | Moose Creek | N/A | 32 |
| | 34 | Browning | 61 | 105 |
| | 61 | Antler | N/A | N/A |
| | 64 | Brock | 54 | 89 |
| | 65 | Tecumseh | 51 | 106 |
| 1B | 91 | Maryfield | 38 | 86 |
| | 122 | Martin | 95 | 151 |
| | 123 | Silverwood | 22 | 74 |
| | 124 A | Kingsley | N/A | 59 |
| | 124 B | Kingsley | N/A | N/A |
| | 125 A | Chester | 37 | 79 |
| | 125 B | Chester | 41 | 101 |
| | 151 | Rocanville | 30 | 81 |
| | 154 | Elcapo | 27 | 80 |
| | 155 | Wolseley | 43 | 91 |
| 2A | 67 | Weyburn | 40 | 85 |
| | 68 | Brokenshell | 40 | 100 |
| | 97 | Wellington | 47 | 82 |
| 2B | 127 A | Francis | 28 | 67 |
| | 127 B | Francis | 20.4 | 40 |
| | 129 | Bratt's Lake | 36 | 100 |
| | 131 A | Baildon | 44 | 133 |
| | 131 B | Baildon | 45 | 155 |
| | 156 A | Indian Head | 30.4 | 69 |
| | 156 B | Indian Head | 22 | 85 |
| | 159 | Sherwood | 38 | 82 |
| | 160 | Pense | 25 | 80 |
| | 161 | Moose Jaw | 36 | 112 |
| | 162 | Caron | 40 | 129 |
| | 191 | Marquis | 25 | 151 |
| 3ASE | 38 A | Laurier | 40.1 | 98 |
| | 38 B | Laurier | 56 | 98 |
| | 39 | The Gap | 40 | 102 |
| 3ASW | 10 | Happy Valley | 28 | 115 |
| | 12 | Poplar Valley | 32 | 130 |
| | 40 | Bengough | N/A | N/A |
| | 42 | Willow Bunch | 31 | 128 |
| | 43 | Old Post | 21 | 114 |
| | 73 A | Stonehenge | 50.8 | 148 |
| | 73 B | Stonehenge | 46 | 139 |
| 3AN | 101 | Terrell | N/A | N/A |
| | 102 | Lake Johnston | 28.5 | 87 |
| | 103 | Sutton | 30 | 98 |
| | 132 A | Hillsborough | 29.5 | 114 |
| | 132 B | Hillsborough | 30 | 134 |
| | 134 | Shamrock | N/A | 17 |
| | 193 A | Eyebrow | 23 | 105 |
| | 193 B | Eyebrow | 45 | 153 |
| 3BS | 17 | Val Marie | 27 | 125 |
| | 18 | Lone Tree | 10 | 145 |
| | 75 | Pinto Creek | 54 | 160 |
| | 76 | Auvergne | N/A | 68 |
| | 77 | Wise Creek | 22 | 173 |
| | 78 | Grassy Creek | 27.5 | 127 |
| | 105 | Glenbain | 29 | 149 |
| | 106 | Whiska Creek | 23 | 86 |
| | 107 | Lac Pelletier | 24 | 98 |
| | 108 | Bone Creek | 30 | 113 |
| 3BN | 138 A | Webb | 35 | 199 |
| | 138 B | Webb | 22 | 136 |
| | 166 | Excelsior | 39 | [illegible] |
| | 167 | Sask. Landing | 53 | [illegible] |
| | 168 A | Riverside | 32 | 152 |
| | 168 B | Riverside | 27.5 | 75 |
| | 226 | Victory | 44 | 123 |
| | 228 | Lacadena | 68 | 153 |
| | 257 | Monet | 57 | 150 |

| Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr |
|---|---|---|---|---|
| 4A | 49 | White Valley | 11 | 102 |
| | 51 | Reno | 15.2 | 107 |
| | 79 | Arlington | 26 | 131 |
| | 109 A | Carmichael | 45.5 | 107 |
| | 109 B | Carmichael | N/A | 216 |
| | 110 | Piapot | 12 | 125 |
| | 111 | Maple Creek | N/A | 106 |
| 4B | 139 | Gull Lake | 28 | 140 |
| | 142 | Enterprise | 28 | 105 |
| | 231 | Happyland | 53 | 112 |
| 5A | 183 | Fertile Belt | 44 | 115 |
| | 211 | Churchbridge | 14 | 71 |
| | 213 | Saltcoats | 16 | 48 |
| | 241 | Calder | 24 | 40 |
| | 243 | Wallace | 20 | 59 |
| | 244 | Orkney | 16 | 50 |
| | 245 A | Garry | 21 | 80 |
| | 245 B | Garry | 25 | 59 |
| | 245 C | Garry | 10 | 43 |
| | 246 | Ituna Bon Accord | 18 | 38 |
| | 247 | Kellross | 6 | 54 |
| | 248 | Touchwood | 12 | 52 |
| 5B | 271 | Cote | 16 | 36 |
| | 273 | Sliding Hills | 5 | 72 |
| | 277 | Emerald | 4 | 48 |
| | 305 | Invermay | 13 | 46 |
| | 307 | Ellros | 6 | 71 |
| | 308 A | Big Quill | 5 | 44 |
| | 308 B | Big Quill | 1 | 56 |
| | 331 | Livingston | 7 | 74 |
| | 336 | Sasman | 4 | 35 |
| | 337 | Lakeview | N/A | 43 |
| | 338 | Lakeside | 1 | 55 |
| | 366 | Kelvington | 1 | 33 |
| | 367 | Ponass Lake | N/A | 65 |
| 6A | 190 A | Dufferin | 29 | 124 |
| | 190 B | Dufferin | 25 | 122 |
| | 190 C | Dufferin | 25 | 110 |
| | 190 D | Dufferin | 36 | 80 |
| | 219 A | Longlaketon | N/A | 43 |
| | 219 B | Longlaketon | 28 | 97 |
| | 220 | McKillop | 27 | 66 |
| | 221 A | Sarnia | 46.7 | 116 |
| | 221 B | Sarnia | 17.5 | 80 |
| | 222 | Craik | 30 | 84 |
| | 251 | Big Arm | 22.5 | 67 |
| | 252 | Arm River | 20 | 66 |
| | 279 | Mount Hope | 14 | 48 |
| | 282 | McCraney | 13 | 59 |
| | 312 | Morris | 23 | 60 |
| | 313 | Lost River | N/A | N/A |
| | 339 | Leroy | 5.4 | 51 |
| | 340 | Wolverine | 19 | 44 |
| | 341 | Viscount | 22 | 54 |
| | 343 | Blucher | 5 | 32 |
| 6B | 223 A | Huron | 37 | 87 |
| | 223 B | Huron | 25 | 104 |
| | 284 A | Rudy | 5 | 51 |
| | 285 B | Rudy | N/A | 51 |
| | 285 A | Fertile Valley | 33 | 91 |
| | 285 B | Fertile Valley | 7.5 | 58 |
| | 286 | Milden | 28 | 97 |
| | 314 | Dundurn | 4 | 37 |
| | 344 | Corman Park | 10 | 43 |
| | 346 | Perdue | 10 | 66 |
| | 376 | Eagle Creek | 8 | 32 |
| | 403 | Rosthern | 10 | 27 |

| Crop Dist. | R.M. No. | Name | Past Week | Since 1-Apr |
|---|---|---|---|---|
| 7A | 287 | St. Andrews | 28 | 76 |
| | 288 | Pleasant Valley | 24 | 64 |
| | 290 A | Kindersley | N/A | 22 |
| | 290 B | Kindersley | 10 | 49 |
| | 290 C | Kindersley | 35 | 44 |
| | 292 | Milton | 31 | 122 |
| | 317 A | Marriott | 32 | 74 |
| | 317 B | Marriott | 19 | 50 |
| | 318 | Mountain View | 31 | 65 |
| | 320 A | Oakdale | 28 | 54 |
| | 320 B | Oakdale | 29 | 52 |
| | 321 | Prairiedale | 32 | 80 |
| 7B | 347 | Biggar | 14 | 30 |
| | 350 A | Mariposa | 50.8 | 67 |
| | 350 B | Mariposa | 19 | 42 |
| | 351 | Progress | 31 | 65 |
| | 352 | Heart's Hill | 14 | 80 |
| | 377 | Glenside | 15 | 30 |
| | 378 | Rosemount | 5 | 34 |
| | 379 | Reford | 13 | 33 |
| | 381 | Grass Lake | 32 | 63 |
| | 382 | Eye Hill | 25.4 | 76 |
| | 409 | Buffalo | 22 | 46 |
| | 410 | Round Valley | 43 | 83 |
| 8A | 395 | Porcupine | N/A | 46 |
| | 397 | Barrier Valley | 3.8 | 54 |
| | 428 | Star City | 3 | 67 |
| | 456 | Arborfield | 5 | 80 |
| | 457 | Connaught | 5 | 71 |
| | 486 | Moose Range | N/A | 63 |
| | 487 | Nipawin | 10 | 106 |
| 8B | 369 | St. Peter | 12 | 28 |
| | 370 A | Humboldt | 6 | 37 |
| | 370 B | Humboldt | 14 | 40 |
| | 371 | Bayne | N/A | 22 |
| | 372 | Grant | N/A | 29 |
| | 400 | Three Lakes | 14 | 54 |
| | 402 | Fish Creek | 31 | 53 |
| | [illegible] | Flett's Springs | 4 | 34 |
| | [illegible] | Kinistino | 2 | 34 |
| | 488 | Birch Hills | 4.3 | 27 |
| 9AE | 488 | Torch River | 10 | 66 |
| | 520 | Paddockwood | 15 | 44 |
| | 521 | Lakeland | 15 | 44 |
| 9AW | 406 | Mayfield | 24 | 36 |
| | 436 | Redberry | 14 | 43 |
| | 438 | Douglas | 12 | 20 |
| | 463 | Duck Lake | 4 | 18 |
| | 467 A | Round Hill | 14 | 48 |
| | 467 B | Round Hill | 16 | 54 |
| | 494 | Canwood | 2 | 27 |
| 9B | 438 | Battle River | 8 | 39 |
| | 440 | Hillsdale | 22.5 | 43 |
| | 442 | Manitou Lake | 33.5 | 48 |
| | 498 A | Parkdale | 18 | 43 |
| | 498 B | Parkdale | 15 | 15 |
| | 499 A | Mervin | 15 | 68 |
| | 499 B | Mervin | N/A | N/A |
| | 501 A | Frenchman Butte | 15 | 36 |
| | 501 B | Frenchman Butte | 29 | 67 |
| | 501 C | Frenchman Butte | 37 | 76 |
| | 502 | Britannia | 47 | 47 |
| | 561 | Loon Lake | 25 | 57 |
| | 588 A | Meadow Lake | NIL | 23 |
| | 588 B | Meadow Lake | 22 | 41 |
| | 622 | Beaver River | N/A | 1 |

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

**Municipality No: A, B, C and D - more than one reporter**

# Cumulative Rainfall

## from April 1 to May 30, 2016




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200
Kilometers
Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture June 1, 2016

© 2016 Government of Saskatchewan

# Cropland Topsoil Moisture Conditions

## May 30, 2016




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture June 1, 2016

# Hay and Pasture Topsoil Moisture Conditions

## May 30, 2016




NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture June 1, 2016

© 2016 Government of Saskatchewan